

Children with Congenital Lip and Palate Splits At I and II Period of Childhood // Annals of the Romanian Society for Cell Biology. – 2021. - Vol. 25. - Issue 4. – P. 1853 – 1858.

6. Kambarova Sh.A. Identification of the morphometric parameters of the cranio-fascial region of children with congenital cleft and palate reflections using a developed research map // Central Asian Journal Of Medical And. – 2021. - Vol. 2. - Issue 3. – P. 286 – 290.
7. Камбаровва С.А. Влияние хирургической манипуляции на морфометрическое развитие лица и челюсти у пациентов с врожденными расстояниями губ и частного двигателя // Новый день в медицине. – 2021. - P. 128 - 130.
8. Kambarova Sh.A., Pulatova Sh.K. Revitalization of nonspecific immunity factors in patients with diffuse phlegmoine of the maxillo facial area using Bakteriofags // New day in medicine. - 2020. - P. 128 - 130.

UDC: 616.314-002-053.2 / 5-07-616.314-74]

THE ROLE OF MINERALIZATION OF MIXED SALIVA IN THE HEALTH OF THE ORAL CAVITY AND THE INFLUENCE OF THE CARIOUSNESS OF THE TEETH

Mirsalihova Firuza Lukmonovna

DSc, docent, Tashkent State Dental Institute,
Department Of children's therapeutic stomatology

E mail: mirsalikhov@gmail.com

Burkhanova Aziza Abduganiyevna

Master of the 1st course at the Department of Children's Therapeutic Dentistry in Tashkent State
Dental Institute.

e-mail: azizakulibaeva@gmail.com

ABSTRACT

The purpose of our study was to study the peculiarities of the biophysical properties and the mineralizing potential of the oral fluid in children during the period of destructing permanent teeth.

Mixed saliva performs a mineralizing function only under the condition of neutral or weakly alkaline pH value when it is oversaturated with calcium and phosphorus ions. Therefore, the physical properties of mixed saliva have a significant impact on the homeostasis of solid teeth tissues. In modern literature, work devoted to the study of the mineralizing potential and biophysical characteristics of saliva in children, contradictory.

The obtained data on the violation of the mineralizing composition and biophysical properties of the oral fluid during the period of destructing permanent teeth testifies to the appropriateness of the appointment of these ages of both the ex- and endogenous prevention of the caries of the teeth immediately after their rubberization.

Keywords: children, dental caries, saliva, mineralization, enamel resistance test.

АННОТАЦИЯ

Целью нашего исследования явилось изучение особенностей биофизических свойств и минерализующего потенциала ротовой жидкости у детей в период прорезывания постоянных зубов.

Смешанная слюна выполняет минерализующую функцию только при условии нейтрального или слабощелочного значения рН, когда она перенасыщена ионами кальция и фосфора. Поэтому физические свойства смешанной слюны оказывают существенное влияние на гомеостаз твердых тканей зубов. В современной литературе работы, посвященные изучению минерализующего потенциала и биофизических характеристик слюны у детей, достаточно противоречивы.

Полученные данные о нарушении минерализующего состава и биофизических свойств ротовой жидкости в период прорезывания постоянных зубов свидетельствует о целесообразности назначения детям данного возраста как экзо-, так и эндогенной профилактики кариеса зубов сразу после их прорезывания.

Ключевые слова: дети, кариес, слюна, минерализация, Тест сопротивления эмали

Relevance. Regarding to modern concepts carious process develops as a result of a complex interaction of general and local factors, which are implemented in the system "organism - saliva - enamel structure" [2,4,8,9]. Among the strong local risk factors for caries, cariogenic microflora in addition, include violation of the composition and properties of oral liquid. Mineralizing properties of oral fluid determined resistance caries dental hard tissues after their eruption, providing the enamel maturation process [2,5,8,9]. Mixed saliva performs mineralizing function only if a neutral or slightly alkaline pH, when it is oversaturated with calcium and phosphorus ions. Respectively, the physical properties of mixed saliva considerable affect the homeostasis of hard tooth tissue [2,8,9]. However, in modern scientific works devoted to the study of potential mineralizing and biophysical characteristics of saliva in children is controversial [2,3,6,7], and are very limited in the Republic of Uzbekistan.

Purpose of the study. To characterize the biophysical properties and mineralizing potential oral fluid in children during a dentition.

Material and methods. To achieve the objectives there were examined 90 children of school ages in the primary school number 42 situated Shaykhantahur district of Tashkent. Children age 7 to 12 years. This age period was chosen as the most vulnerable to dental caries, in the period of mixed dentition and dentition.

We have carried out inspection of the features of the biophysical properties of the mixed saliva in this age: the pH and buffering capacity of the oral fluid, the viscosity and the rate of secretion was studied using a test kit «Saliva-check Buffer» (GC).

Mineralizing potential of mixed saliva (MPU) has been studied by the method of PA Leus (1977). Caries tooth enamel has been studied using the FER test method VR Okushko, LI Kosarev IK Lutsk (1983). Determination mineralizing potential of saliva (MPU)

The material for crystallographic studies was taken from the oral fluid of school pupils, which was collected on an empty stomach. Oral liquid was applied on a glass slide in a drop of 1 ml 18-25 °C and dried at a temperature in the horizontal position. Investigation of the structure of saliva samples was performed using an optical microscope.

Results and discussion. The evaluation results of the study mineralizing potential of saliva (MPS) carried out by the method of PA Leus (1977) estimate the character drawing.

1 ball-placers randomly distributed irregularly shaped structures

2 points - a fine line grid over the entire field of view

3 points, the individual crystals of irregular shape on a grid background and clumps

4 points - a medium-sized tree-like crystals

5 points - clearly large, like karkon fern or crystal structure.

MPS assessed value on the following parameters:

0-1 3,1-4,0 very low high

1.1-2.0 low 4, -5 very high

2,1-3 satisfactory

MPS (PA Leus, 1977) also gives an indication of the oral fluid saturation of trace elements [4.12].

Sets the type of microcrystallisation in oral fluid. 1-type matched saliva rich in trace elements and is an elongated prismatic crystal structures, often with radial orientation. 2 type - consistent with the average level of micronutrients saturated saliva and looked like isometric located fragments of crystals without a clear orientation.

3-type matched saliva oral fluid poor micronutrients - scattered small isolated fragments of non-oriented crystals.

The children also performed a standard dental examination, assess the state of hard tissue of teeth, etc. with enamel resistance test (TER), and mineralized potential of saliva (IPU).

TER - Test (Okushko VR, LI Kosarev, Lutsk IK, 1983) evaluated the stability of tooth enamel to acid attack. Therefore, the method of the central incisor of the upper jaw was purified from the soft plaque 1% H₂O₂ solution was dried with a dry cotton swab. In the middle of the vestibular surface of the tool for 2-3 seconds pipetted mordant acid 1.5-2.5 mm in diameter. Etchant then removes dry cotton swab and stained with 2% methylene blue solution. Dye rented a dry cotton swab the deleting movements, completely taking off the enamel surface. Etched portion is painted in blue color of varying intensity. To evaluate the intensity of staining was used a 10-point scale typographical tint of blue from 1.0 to 10.0 points color intensity.

If the color intensity from 1.0 to 3.0 children surveyed are at high caries (HCR) group, painting 4-5 moderate (RBM), the color of 6-7 points lower (LCR), the color of more than 8 points - very low (VLCR), the maximum risk of caries.

In assessing the results of the study FER test set that out of 90 students surveyed in only 18 (20%) of children set a high level of KR tooth enamel (WRC), the remaining 82 (80.0%) children had a different level of the CD. So in 24 (26.7%) patients were identified moderate KR (RBM) in 27 (30.0%) lower the Kyrgyz Republic (LCR) and 21 (23.3%) children is very low Raman (VLCR) (Table1).

Table 1

Index of TER-test

| Number of examined children 90 | HCR | | NCR | | LCR | | VLCR | |
|-----------------------------------|-----|------|-----|------|-----|----|------|------|
| | abs | % | abs | % | abs | % | abs | % |
| | 18 | 20,0 | 21 | 23,3 | 27 | 30 | 24 | 26,7 |

Analysis of the data table shows that the majority of children studied (56.7%) have a low or very low resistance of teeth to acid exposure that forecasts about the development of multiple dental caries. When the study results MPS study in primary school children found that in children with SRS tooth enamel in 50.0% of cases had a high level of MEA and 11.1% had a very high level of the IPU, the remaining 38.9% satisfactory. Children with low or very low MPS absent.

In the study of the Ministry of Railways in the second RBM children, as shown by the table number 2 was dominated by students with mild MPS level - 71.4%, the lowest MPS is set at 19.0%, the highest level was observed in only 9.9%. Students with very high and very low levels of MPS no.

Children with LCR tooth enamel in 51.8% of cases diagnosed MPS low in 25.9% of cases satisfactory. In 22.2% of cases registered with the Ministry of Railways is very low. High or very high MPS is detected.

Table2

The frequency of the level of the IPU in children, depending on the tooth enamel caries

| Index of TER | 1 group, n=18 HCR | | 2 group, n=21 NCR | | 3 group, n=27 LCR | | 4 group, n=24 VLCR | |
|---------------------|----------------------|------|----------------------|------|----------------------|------|-----------------------|------|
| | abs | % | abs | % | abs | % | abs | % |
| Level MPS | - | - | - | - | 6 | 22,2 | 9 | 37,5 |
| Very low, 0-1 | - | - | 4 | 19,0 | 14 | 51,8 | 12 | 50,0 |
| Low, 1,1-2 | 7 | 38,9 | 15 | 71,4 | 7 | 25,9 | 3 | 12,5 |
| Satisfactory, 2,1-3 | 9 | 50,0 | 2 | 9,9 | - | - | - | - |
| High, 3,1-4 | 2 | 11,1 | - | - | - | - | - | - |
| Very high, 4,1-5 | | | | | | | | |

In the fourth group of children with enamel, VLCR established the predominance of low and very low MPS respectively 50.0% and 37.5%. Satisfactory MPS 12.5% of cases. High and very high MPS completely absent.

Thus, the analysis of the data indicates a violation of mineralizing function of oral fluid in children of primary school age, expressed decrease mineralizing potential of saliva at the LCR and VLCR tooth enamel.

It revealed a direct correlation relationship between the studied parameters of CR tooth enamel and the IPU. Since the first lower productivity promotes the deterioration of the second and they are interdependent.

Studying of mineralizing potential of oral fluid in primary school children also has substantial diagnostic value for identifying the development opportunities of cariogenic situation in the oral cavity in primary school children during the period of the dentition.

The oral liquid is one of informative biological fluids in the human body. It plays a significant role in the regulation of homeostasis of the oral cavity. During the study, various parameters of the oral liquid can be used as a marker of a number of diseases of the oral cavity.

Therefore, the diagnostic value of MPS oral liquid children is unquestionable. In accordance with the received data, it is advisable to carry out a test control mineralizing potential of mixed saliva to predict possible development of cariogenic situation that contributes to the early diagnosis of the establishment of increased cariogenic situation and timely appointment of prophylactics.

Conclusion. It was determined the four degree of resistance to tooth decay HCR, NCR, LCR VLCR. The changes of the biophysical properties of mixed saliva: decrease of pH, buffer capacity and rate of saliva secretion in the LCR and VLCR tooth enamel in children which is a forecast of possible development of an active flow of caries during the eruption of permanent teeth in children and appropriateness of the destination as the exo and endogenous caries prevention immediately after the eruption.

Литература / References

1. Borovsky S.V. oral biology, V.E. Borovsky, V.K. Leontiev. Moscow: Medicine. -2001.-304.
2. Denisov A.B. Diagnostic informativeness saliva. Prospects /A.B. research. Denisov // Dental Review. Education, science and practice of dentistry. Abstracts of the All-Russian Dental Forum. -Moscow, 2004. C. 253-256.
3. Leontiev V.K. Dental Caries - a disease of civilization / V.K. Leontiev // biosphere. -2010.-№ 3.-S. 392-396.
4. Leontiev E.Y. et al. remineralization therapy using TOOTN MAUSSE and Mi Pastr plus (GC) // Problems of dentistry. -2012.-№ 1.-S.33-36.
5. Leus P.A. Clinical and experimental study of pathogenesis pathogenetic conservative treatment and prevention of dental caries. Author. Dis. Dr. Med. -M.1977.-30.
6. Nowicka I.K. The correlation between the intensity of caries and indicators mineralizing fluid capacity and infants // of Jornal of Health Sciencas.-2014.-Vol.04.-N 03.-P.011-020.
7. Savushkina N.A., Kobiyasova I.V. Features mineralized saliva function in adolescents puberty and methods of its correction // Dentistry childhood and prevention. - 2003. - № 3-4. - S.28-32.

8. Preethi B.P., Pyati A., Dodawad R. Evaluation of flow rate, pH, buffering capacity, calcium, total protein and total antioxidant levels of saliva in caries free and caries active children // Biomed, Research. - 2010.- Vol.21, N 3. - P.289-294.

9. Serum and salivary minerals in dental caries / M.Jawed, S.M. Shahid, A. Rehman et al. // J. Dow. University of Health Sciences. - 2009. - Vol.3, N 2. - P. 61-65.

УДК: 617.52/53-001.16-089.844 : 611.018

**МОРФОЛОГИЧЕСКОЕ ОБОСНОВАНИЕ ВОЗМОЖНОСТИ ПРИМЕНЕНИЯ
ИНТРАОПЕРАЦИОННО РАСТЯНУТЫХ ТКАНЕЙ ПРИ ПЛАСТИКЕ РУБЦОВЫХ
ДЕФОРМАЦИЙ ЛИЦА И ШЕИ**

Тешабоев М.Г.

Андижанский государственный медицинский институт

teshaboev2013@mail.ru

В данной статье представлены обоснованные рекомендации по применению интраоперационного баллонного растяжения тканей для устранения рубцовых деформаций последствиями ожогов лица и шеи. Полученные морфологические результаты показывают, что метод интраоперационного баллонного растяжения тканей обеспечивает жизнеспособность лоскута.

Вывод. Таким образом, интраоперационное растягивание мягких тканей по разработанной в клинике схеме не вызывает патологические изменения общей архитектоники кожи, не нарушает микрорельеф поверхности и сетчатую структуру волоконной основы кожи.

Ключевые слова: интраоперационные баллонные растяжение ткани, рубцовая деформация последствиями ожогов, лицо, шея, лоскут.

**MORPHOLOGICAL SUBSTANTIATION OF THE POSSIBILITY OF APPLICATION OF
INTRAOPERATIVELY STRETCHED TISSUES IN PLASTIC OF FACE AND NECK
SCAR DEFORMATIONS**

Teshaboev M.G.

Andijan State medical institute

teshaboev2013@mail.ru

This article presents reasonable recommendations on the use of intraoperative balloon stretching of tissues to eliminate cicatricial deformities resulting from burns to the face and neck. The obtained morphological results show that the method of intraoperative balloon stretching of tissues ensures the viability of the flap.

Output. Thus, intraoperative stretching of soft tissues according to the scheme developed in the clinic does not cause pathological changes in the general architectonics of the skin, does not violate the surface microrelief and the reticular structure of the fiber base of the skin.

Key words: intraoperative balloon tissue stretching, cicatrice deformity caused by burns, face, neck, flap.

**ЮЗ ВА БЎЙИН КУЙИК АСОРАТИ БОРЛАРНИ ИНТРАОПЕРАЦИОН БАЛЛОН
ТЎҚИМА ЧЎЗИШ ПЛАСТИКАСИНИ МОРФОЛОГИК АСОСЛАШ**

Тешабоев М.Г.